

CLAIMS

Add
B'

1. Products containing a non-depleting CD4 monoclonal antibody and a non-depleting CD8 monoclonal antibody as a combined preparation for simultaneous, separate or sequential use in inducing tolerance to an antigen.
2. Products according to Claim 1, further containing a depleting CD4 monoclonal antibody and/or a depleting CD8 monoclonal antibody.
3. A non-depleting CD4 monoclonal antibody for use in a method of treatment of the human or animal body by surgery or therapy.
4. A non-depleting CD8 monoclonal antibody for use in a method of treatment of the human or animal body by surgery or therapy.
5. An antibody according to Claim 3 or 4 for use in inducing tolerance to an antigen.
6. Use of a non-depleting CD4 monoclonal antibody in the preparation of a medicament for use in inducing tolerance to an antigen.
7. Use of a non-depleting CD8 monoclonal antibody in the preparation of a medicament for use in inducing tolerance to an antigen.
8. An agent for use in inducing tolerance to an antigen, which agent comprises a non-depleting CD4 monoclonal antibody and a non-depleting CD8 monoclonal antibody.
9. An agent according to Claim 8, further comprising a depleting CD4 monoclonal antibody and/or a depleting CD8 monoclonal antibody.
10. A pack comprising as separate components a non-depleting CD4 monoclonal antibody and a non-depleting CD8 monoclonal antibody.

- 44 -

11. A pack according to Claim 10, further comprising as separate components a depleting CD4 monoclonal antibody and/or a depleting CD8 monoclonal antibody.

5 12. Non-depleting CD4 and CD8 monoclonal antibodies for use together in a method of treatment of the human or animal body by surgery or therapy.

13. Antibodies according to Claim 12, for use together in inducing tolerance to an antigen.

10 14. Antibodies according to Claim 12 or 13, further comprising a depleting CD4 monoclonal antibody and/or a depleting CD8 monoclonal antibody.

15 15. A method of conferring on a subject tolerance to an antigen it is wished to administer thereto, which method comprises administering to the subject an effective amount of a non-depleting CD4 monoclonal antibody and a non-depleting CD8 monoclonal antibody and of the antigen. ✓

20 16. A method of conferring on a subject tolerance to an antigen already possessed by the subject, which method comprises administering to the subject an effective amount of a non-depleting CD4 monoclonal antibody and a non-depleting CD8 monoclonal antibody. ✓

25 17. A method according to Claim 15 or 16, further comprising administering to the subject an effective amount of a depleting CD4 monoclonal antibody and/or a depleting CD8 monoclonal antibody prior to administration of the non-depleting CD4 and CD8 monoclonal antibodies.

add
B1

add
D4